

International Standard



6800

INTERNATIONAL ORGANIZATION FOR STANDARDIZATION • МЕЖДУНАРОДНАЯ ОРГАНИЗАЦИЯ ПО СТАНДАРТИЗАЦИИ • ORGANISATION INTERNATIONALE DE NORMALISATION

Animal and vegetable fats and oils — Determination of the composition of fatty acids in the 2-position

Corps gras d'origines animale et végétale — Détermination de la composition des acides gras en position 2

First edition — 1985-12-15

STANDARDSISO.COM : Click to view the full PDF of ISO 6800:1985

UDC 665.2/.3:543.8

Ref. No. ISO 6800-1985 (E)

Descriptors : agricultural products, animal fats, vegetable fats, animal oils, vegetable oils, chemical analysis, determination of content, fatty acids.

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work.

Draft International Standards adopted by the technical committees are circulated to the member bodies for approval before their acceptance as International Standards by the ISO Council. They are approved in accordance with ISO procedures requiring at least 75 % approval by the member bodies voting.

International Standard ISO 6800 was prepared by Technical Committee ISO/TC 34, *Agricultural food products*.

Users should note that all International Standards undergo revision from time to time and that any reference made herein to any other International Standard implies its latest edition, unless otherwise stated.

Animal and vegetable fats and oils — Determination of the composition of fatty acids in the 2-position

1 Scope

This International Standard specifies a method for the determination of the composition of fatty acids which are in the 2-position (β or internal position) of glyceride molecules in animal and vegetable fats and oils.

2 Field of application

Because of the nature of pancreatic lipase action, the method is applicable only to fats and oils with a melting point below 45 °C.

The method is not unreservedly applicable to all fats and oils, particularly those containing substantial amounts of

- fatty acids with 12 or fewer carbon atoms (copra oil, palm kernel oil, butyric butter fats);
- fatty acids with 20 and more carbon atoms and of a high degree of unsaturation (more than four double bonds) (fish oil and marine animal oil);
- fatty acids which have secondary groups containing oxygen.

3 References

ISO 660, *Animal and vegetable fats and oils — Determination of acid value and of acidity.*

ISO 661, *Animal and vegetable fats and oils — Preparation of test sample.*

ISO 5508, *Animal and vegetable fats and oils — Analysis by gas-liquid chromatography of methyl esters of fatty acids.*

ISO 5509, *Animal and vegetable fats and oils — Preparation of methyl esters of fatty acids.*

ISO 5555, *Animal and vegetable fats and oils — Sampling.*

4 Principle

After neutralization, where necessary, of any free fatty acids, purification of the test portion by column chromatography. Partial enzymatic hydrolysis of the glycerides to yield 2-monoglycerides. Separation of the monoglycerides by thin layer chromatography and determination of their fatty acid composition by gas chromatography.

5 Reagents

All reagents shall be of recognized analytical quality and the water used shall be distilled water or water of at least equivalent quality.

5.1 Reagents for the purification of the test portion

5.1.1 2-propanol, or ethanol 95 % (V/V).

5.1.2 Hexane (if available) or light petroleum (boiling range 30 to 60 °C).

5.1.3 2-propanol 50 % (V/V) or ethanol 50 % (V/V).

5.1.4 Sodium hydroxide solution, 0,5 mol/l.

5.1.5 Phenolphthalein solution, 1 g per 100 ml of ethanol 95 % (V/V).

5.1.6 Activated neutral alumina, for chromatography, Brockmann activity I, recently activated for 2 h at 260 °C and kept in a desiccator.

5.1.7 Nitrogen.

5.2 Reagents for hydrolysis of the triglycerides

5.2.1 **Diethyl ether**, free from peroxides.

5.2.2 **Hydrochloric acid**, 6 mol/l.

5.2.3 **Sodium cholate** solution, 1 g/l, of enzymatic quality.

5.2.4 **Calcium chloride** (CaCl_2) solution, 220 g/l.

5.2.5 **Buffer solution**, 2-amino-2-(hydroxymethyl) propane-1,3-diol¹⁾, 1 mol/l, adjusted to pH 8 with 6 mol/l hydrochloric acid (5.2.2) using a pH-meter.

Store this solution at 0–4 °C and use within 14 days.

5.2.6 **Pancreatic lipase**, with an activity of between 8 and 20 units/mg (see the annex).

Store dry in a refrigerator. Before use, bring a portion of the powder to ambient temperature.

NOTE — Lipase of suitable activity is available commercially. If preferred, the lipase may be prepared and assayed in accordance with the procedure described in the annex.

5.3 Reagents for the isolation of the 2-monoglycerides

5.3.1 **Ethanol**.

5.3.2 **Hexane** (if available), or **light petroleum** (boiling range 30 to 60 °C).

5.3.3 **Acetone**.

5.3.4 **Silica powder**, with binder, for thin layer chromatography.

5.3.5 **Developing solvent**, prepared as follows :

hexane (if available) or light petroleum :	70 ml
diethyl ether :	30 ml
formic acid, minimum 98 % (V/V) :	1 ml

5.3.6 **2', 7'-Dichlorofluorescein**, indicator solution, 2 g per litre of methanol, rendered slightly alkaline by addition of a drop of 1 mol/l sodium hydroxide per 100 ml of the solution.

5.4 Reagents for the analysis of the 2-monoglycerides by gas chromatography

See ISO 5508 and ISO 5509.

6 Apparatus

Usual laboratory equipment, and in particular

6.1 Apparatus for the purification of the test portion

6.1.1 **Water-bath**, thermostatically controlled, and capable of being maintained at 30 to 40 °C.

6.1.2 **Glass column**, for chromatography, 13 mm internal diameter and 400 mm in length, equipped with a sintered glass plate and a tap.

6.1.3 **Rotary evaporator**, with 250 ml flask.

6.1.4 **Tubing**, for nitrogen bubbling.

6.1.5 **Separating funnel**, 500 ml capacity.

6.1.6 **Round-bottom flask**, 100 ml capacity.

6.2 Apparatus for the hydrolysis of the triglycerides

6.2.1 **Centrifuge** (if necessary).

6.2.2 **Centrifuge tube**, 10 ml capacity, with ground stopper.

6.2.3 **Vibrating electric shaker**, for vigorous agitation of the centrifuge tube.

6.2.4 **Water-bath**, thermostatically controlled, and capable of being maintained at $40 \pm 0,5$ °C.

6.2.5 **Hypodermic syringe**, 1 ml capacity, with thin needle.

6.2.6 **Stop-watch**.

6.3 Apparatus for the isolation of the 2-monoglycerides

6.3.1 **Developing tank**, for thin layer chromatography, with ground glass lid, suitable for containing glass plates 200 mm × 200 mm.

6.3.2 **Spreader and rack**, for preparation of the plates.

6.3.3 **Glass plates**, 200 mm × 200 mm.

6.3.4 **Microsyringe**, capable of dispensing drops of 3 to 4 µl as a continuous uniform band.

1) Alternative names : tris-(hydroxymethyl) methylamine; tris-(hydroxymethyl) aminomethane.

6.3.5 Apparatus for spraying the indicator solution on to the plates.

6.3.6 Microspatula.

6.3.7 Oven, capable of being maintained at $103 \pm 2^\circ\text{C}$.

6.3.8 Ultraviolet lamp, for examining chromatographic plates, for example with a wavelength of 254 nm.

6.3.9 Round-bottom flask, 25 ml capacity, **with air condenser** of approximately 1 m length with ground joint.

6.3.10 Conical flask, 250 ml capacity, with ground stopper.

6.3.11 Conical flask, 50 ml (if necessary).

6.3.12 Filter, sintered glass, porosity P 40 (16 to 40 μm) (if necessary).

6.3.13 Desiccator.

6.4 Apparatus for the analysis of the 2-monoglycerides by gas chromatography

See ISO 5508 and ISO 5509.

7 Sampling

See ISO 5555.

8 Procedure

8.1 Preparation of test sample

Prepare the test sample from the laboratory sample in accordance with ISO 661.

8.2 Determination of the acidity of the test sample

Determine the acidity of the test sample in accordance with ISO 660.

If the acidity is below 3 % (m/m), purify the sample through alumina in accordance with 8.4.

If the acidity exceeds 3 % (m/m), first neutralize the sample with sodium hydroxide in the presence of solvent in accordance with 8.3, then purify through alumina in accordance with 8.4.

8.3 Neutralization with sodium hydroxide

Dissolve about 10 g of the test sample in 100 ml of hexane (if available) or light petroleum (5.1.2) and transfer the solution to the separating funnel (6.1.5). Add 50 ml of 2-propanol or ethanol 95 % (V/V) (5.1.1), a few drops of phenolphthalein solution (5.1.5) and a volume of sodium hydroxide solution

(5.1.4) equivalent to the free acidity of the fat or oil, with an excess of 0,5 %. Shake vigorously for 1 min, add 50 ml of water, shake again and allow to stand. When separated, run off the lower layer containing the soap and any intermediate layers (mucilages and insoluble substances). Wash the hexane or light petroleum solution of neutralized oil with successive portions of 25 or 30 ml of a solution of 2-propanol or ethanol (5.1.3) until the pink colour of the phenolphthalein disappears.

Transfer the solution into the rotary evaporator flask (6.1.3) and remove most of the solvent by evaporating under reduced pressure. Dry the oil at 30 to 40 $^\circ\text{C}$ under reduced pressure using a nitrogen stream (5.1.7) until the solvent is completely removed.

8.4 Purification of the test portion through alumina

Prepare a suspension of 15 g of activated alumina (5.1.6) in 50 ml of hexane (if available) or light petroleum (5.1.2) and pour, while shaking, into a glass column for chromatography (6.1.2). Ensure that the alumina settles evenly and allow the solvent level to fall to 1 to 2 mm above the upper level of the adsorbent. Carefully pour into the column a solution prepared by dissolving a test portion of 5 g, neutralized if necessary, in 25 ml of hexane (if available) or light petroleum (5.1.2) and collect all the liquid which elutes from the column in a 100 ml round-bottom flask (6.1.6).

Remove most of the solvent by evaporation under reduced pressure, then dry the oil at 30 to 40 $^\circ\text{C}$ using a nitrogen stream (5.1.7) until the solvent is completely removed.

8.5 Hydrolysis of the triglycerides

Weigh approximately 0,1 g of the purified test portion (8.4) in a 10 ml centrifuge tube (6.2.2). If this test portion is solid, place the tube in a water-bath at 60–65 $^\circ\text{C}$ until it has liquefied or for not longer than 40 s, even if by then the test portion is not completely liquid. Bring the temperature of the tube to about 40 $^\circ\text{C}$ and continue the procedure without delay.

Place the centrifuge tube containing the purified test portion in the water-bath at $40 \pm 0,5^\circ\text{C}$. Add 2 ml of the buffer solution (5.2.5), 0,5 ml of sodium cholate solution (5.2.3) and 0,2 ml of calcium chloride solution (5.2.4) and shake carefully. Add 20 mg of lipase (5.2.6), insert the stopper and shake manually for exactly 1 min with the tube in the water-bath.

Remove the tube from the water-bath and shake vigorously for exactly 2 min using the shaker (6.2.3).

Add 1 ml of hydrochloric acid (5.2.2) and 1 ml of diethyl ether (5.2.1). Insert the stopper and shake vigorously using the shaker (6.2.3). Centrifuge and transfer the organic phase to a test-tube using the syringe (6.2.5). If the test portion was solid at ambient temperature, repeat the extraction with a further 1 ml portion of diethyl ether and combine the extracts in the test-tube.